

Listing of Claims:

This listing of claims will replace all prior versions, and listings, of claims in the application:

1. (Currently Amended) A method for treating proliferative skin diseases comprising the administration of an effective amount of 1.0 % tazarotene gel and an effective amount of a corticosteroid[[.]], wherein said corticosteroid is sSelected from the group consisting of fluocinolone acetamide, mometasone furoate, fluocinonide, diflorasone diacetate, fluticasone propionate, betamethasone dipropionate, clobetasol propionate, and betamethasone valerate.
2. (Cancelled) The method of claim 1 wherein said corticosteroid is selected from the group consisting of fluocinolone acetamide, mometasone furoate, fluocinonide, diflorasone diacetate, fluticasone propionate, betamethasone dipropionate, clobetasol propionate, and betamethasone valerate.
3. (Cancelled) The method of claim 1 wherein tazarotene is applied as a 0.1% gel.
4. (Currently Amended) The method of claim 1 wherein said corticosteroid is [[a]] selected from the group consisting of mometasone furoate, fluocinonide, alclometasone dipropionate, and betamethasone valerate.
5. (Original) The method of claim 4 wherein said corticosteroid is selected from the group consisting of alclometasone dipropionate, and betamethasone valerate.
6. (Currently Amended) A method for treating psoriasis in a human subject by topically applying to the psoriatic skin of said subject an effective amount of 1.0 % tazarotene and an effective amount of a corticosteroid.
7. (Cancelled) The method of claim 6 wherein tazarotene is applied as a 0.1% gel.

8. (Cancelled) The method of claim 7 wherein said corticosteroid is a cream.
9. (Cancelled) The method of claim 8 wherein said corticosteroid is selected from the group consisting of mometasone furoate, fluocinonide, alclometasone dipropionate, and betamethasone valerate.
10. (Original) The method of claim [[9]] 6 wherein said corticosteroid is selected from the group consisting of alclometasone dipropionate, and betamethasone valerate.
11. (Currently Amended) The method of claim [[6]] 10 wherein tazarotene is administered once daily in the evening and the corticosteroid is administered once daily in the morning.
12. (New) The method of Claim 6 wherein the 1.0 % tazarotene is applied as a 1.0 % gel.
13. (New) The method of Claim 6 wherein the 1.0 % tazarotene is applied as a 1.0 % cream.
14. (New) The method of Claim 6 wherein the 1 corticosteroid is fluocinonide.